

Mini Nutritional Assessment

MNA[®]

Last name:		First name:		
Sex:	Age:	Weight, kg:	Height, cm:	Date:

Complete the screen by filling in the boxes with the appropriate numbers.
Add the numbers for the screen. If score is 11 or less, continue with the assessment to gain a Malnutrition Indicator Score.

Screening

A Has food intake declined over the past 3 months due to loss of appetite, digestive problems, chewing or swallowing difficulties?
0 = severe decrease in food intake
1 = moderate decrease in food intake
2 = no decrease in food intake

B Weight loss during the last 3 months
0 = weight loss greater than 3kg (6.6lbs)
1 = does not know
2 = weight loss between 1 and 3kg (2.2 and 6.6 lbs)
3 = no weight loss

C Mobility
0 = bed or chair bound
1 = able to get out of bed / chair but does not go out
2 = goes out

D Has suffered psychological stress or acute disease in the past 3 months?
0 = yes 2 = no

E Neuropsychological problems
0 = severe dementia or depression
1 = mild dementia
2 = no psychological problems

F Body Mass Index (BMI) (weight in kg) / (height in m²)
0 = BMI less than 19
1 = BMI 19 to less than 21
2 = BMI 21 to less than 23
3 = BMI 23 or greater

Screening score (subtotal max. 14 points)

12-14 points: Normal nutritional status
8-11 points: At risk of malnutrition
0-7 points: Malnourished

For a more in-depth assessment, continue with questions G-R

Assessment

G Lives independently (not in nursing home or hospital)
1 = yes 0 = no

H Takes more than 3 prescription drugs per day
0 = yes 1 = no

I Pressure sores or skin ulcers
0 = yes 1 = no

J How many full meals does the patient eat daily?
0 = 1 meal
1 = 2 meals
2 = 3 meals

K Selected consumption markers for protein intake

- At least one serving of dairy products (milk, cheese, yoghurt) per day yes no
- Two or more servings of legumes or eggs per week yes no
- Meat, fish or poultry every day yes no

0.0 = if 0 or 1 yes
0.5 = if 2 yes
1.0 = if 3 yes

L Consumes two or more servings of fruit or vegetables per day?
0 = no 1 = yes

M How much fluid (water, juice, coffee, tea, milk...) is consumed per day?
0.0 = less than 3 cups
0.5 = 3 to 5 cups
1.0 = more than 5 cups

N Mode of feeding
0 = unable to eat without assistance
1 = self-fed with some difficulty
2 = self-fed without any problem

O Self view of nutritional status
0 = views self as being malnourished
1 = is uncertain of nutritional state
2 = views self as having no nutritional problem

P In comparison with other people of the same age, how does the patient consider his / her health status?
0.0 = not as good
0.5 = does not know
1.0 = as good
2.0 = better

Q Mid-arm circumference (MAC) in cm
0.0 = MAC less than 21
0.5 = MAC 21 to 22
1.0 = MAC 22 or greater

R Calf circumference (CC) in cm
0 = CC less than 31
1 = CC 31 or greater

Assessment (max. 16 points)

Screening score

Total Assessment (max. 30 points)

Malnutrition Indicator Score

24 to 30 points	<input type="checkbox"/>	Normal nutritional status
17 to 23.5 points	<input type="checkbox"/>	At risk of malnutrition
Less than 17 points	<input type="checkbox"/>	Malnourished

References

- Vellas B, Villars H, Abellan G, et al. Overview of the MNA[®] - Its History and Challenges. *J Nutr Health Aging*. 2006; **10**:456-465.
- Rubenstein LZ, Harker JO, Salva A, Guigoz Y, Vellas B. Screening for Undernutrition in Geriatric Practice: Developing the Short-Form Mini Nutritional Assessment (MNA-SF). *J Gerontol*. 2001; **56A**: M366-377
- Guigoz Y. The Mini-Nutritional Assessment (MNA[®]) Review of the Literature - What does it tell us? *J Nutr Health Aging*. 2006; **10**:466-487.

© Société des Produits Nestlé, S.A., Vevey, Switzerland, Trademark Owners

© Nestlé, 1994, Revision 2009. N67200 12/99 10M

For more information: www.mna-elderly.com